

## Natural Resource Biometrics

### Testing differences between two samples with similar variance

If the F test for variance does not test significantly different you can calculate a pooled variance as:

$$s_p^2 = \frac{df_1 s_1^2 + df_2 s_2^2}{df_1 + df_2}$$

### Testing the difference between two means

The means of two samples can be tested for difference by:

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_p^2}{n_1} + \frac{s_p^2}{n_2}}}$$

### Also See:

Chapter 9 - Significance of a Difference between Two Means pages 108-124 in:

Phillips, J. L. 2000. How to think about statistics. W. H. Freeman and Co. New York. 202 pp. ISBN 0-7167-3654-3

Chapter 9 - Two-Sample Hypotheses pages 126-130 in:

Zar, J. H. 2007. Biostatistical Analysis. Prentice-Hall, Inc. Englewood Cliffs, New Jersey. 718 pp.



Natural Resources Biometrics by David R. Larsen is licensed under a [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).

Author: Dr. David R. Larsen  
Created: October 6, 2013  
Last Updated: August 18, 2014